No.



Hennsylbania Agricultural Experiment Station

MECCENS, THERE HAS BEEN PRESENTED TO THE

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED DISTINCT VARIETY OF SEXUALLY REPRODUCED, OR TUBER PROPAGATED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF TWENTY YEARS FROM THE DATE OF THIS GRANT, SUBJECT 10 THE PAYMENT OF THE REQUIRED FEES AND PERIODIC REPLENISHMENT OF VIABLE BASIC SEED OF THE VARIETY IN A JUBILIC REPOSITORY AS PROVIDED BY LAW, THE IGHT TO EXCLUDE OTHERS FROM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT, OR ORTING IT, OR EXPORTING IT, OR CONDITIONING IT FOR PROPAGATION, OR STOCKING IT FOR ANY OF THE E PURPOSE, OR CONDITIONING IT FOR PROPAGATION, OR STOCKING IT FOR ANY OF THE ABOVE E, OR USING IT IN PRODUCING A HYBRID OR DIFFERENT VARIETY THEREFROM, TO THE EXTENT BY THE PLANT VARIETY PROTECTION ACT. IN THE UNITED STATES SEED OF THIS VARIETY SOLD BY VARIETY NAME ONLY AS A CLASS OF CERTIFIED SEED AND (2) SHALL CONFORM TO THE NUM NERATIONS SPECIFIED BY THE OWNER OF THE RIGHTS, (84 STAT, 1542, AS AMENDED, 7 U.S.C. 2321 ET SEO

BENTGRASS, CREEPING

'Seaside II'

In Testimonn Mercert, I have hereunto set my hand and caused the seal of the Mant Bariety Brotection Office to be affixed at the City of Washington, D.C. this eleventh day of March, in

NAME (Please print or type)

CAPACITY OR TITLE

10/17/96

Associate Dean

STD-470 (03-86) Previous editions are to be destroyed.

Charles R. Krueger

NAME (Please print or type)

CAPACITY OR TITLE

(See reverse for instructions and information collection burden statement)

DATE

INSTRUCTIONS

GENERAL: To be effectively filed with the Plant Variety Protection Office (PVPO), ALL of the following items must be received in PVPO: (1) Completed application form signed by the owner; (2) completed Exhibits A, B, C, E; (3) at least 2,500 viable untreaseds, or for tuber reproduced varieties verification that a viable (in the sense that it will reproduce an entire plant) tissue culture be deposited and maintained in an approved public repository; (4) check drawn on a U.S. bank for \$2,450 (\$300 filing fee \$2,150 examination fee), payable to "Treasurer of the United States" (See Section 97.6 of the Regulations and Rules of Practice Partial applications will be held in the PVPO for not more than 90 days, then returned to the applicant as unfiled. Mail application other requirements to Plant Variety Protection Office, AMS, USDA, Room 500, NAL Building, 10301 Baltimore Blvd., Beltsville, National Corrections on the application form and exhibits must be initialed and dated. DO NOT use masking materials to make corrections a certificate is allowed, you will be requested to send a check payable to "Treasurer of the United States" in the amount of \$300 issuance of the Certificate.

Plant Variety Protection Office Telephone: (301) 504-5518

ITEM

- 16a. Give: (1) the genealogy, including public and commercial varieties, lines, or clones used, and the breeding method;
 - (2) the details of subsequent stages of selection and multiplication;
 - (3) evidence of uniformity and stability; and
 - (4) the type and frequency of variants during reproduction and multiplication and state how these variants may be identified.
- 16b. Give a summary of the variety's distinctness. Clearly state how this application variety may be distinguished from all oth varieties in the same crop. If the new variety is most similar to one variety or a group of related varieties:
 - (1) identify these varieties and state all differences objectively;
 - (2) attach statistical data for characters expressed numerically and demonstrate that these are clear differences;
 - (3) submit, if helpful, seed and plant specimens or photographs (prints) of seed and plant comparisons which clear indicate distinctness.
- 16c. Exhibit C forms are available from the PVPO for most crops; specify crop kind. Fill in Exhibit C (Objective Description of Variety) form as completely as possible to describe your variety.
- Optional additional characteristics and/or photographs. Describe any additional characteristics that cannot be accurately conveyed in Exhibit C. Use comparative varieties as is necessary to reveal more accurately the characteristics that are difficult to describe, such as plant habit, plant color, disease resistance, etc.
- 16e. Section 52(5) of the Act requires applicants to furnish a statement of the basis of the applicant's ownership. An Exhibit E form is available from the PVPO.
- 17. If "Yes" is specified (seed of this variety be sold by variety name only, as a class of certified seed), the applicant may NO reverse this affirmative decision after the variety has been sold and so labelled, the decision published, or the certificate issued. However, if "No" has been specified, the applicant may change the choice. (See Regulations and Rules of Practice Section 97.103).
- 20. See Sections 41, 42, and 43 of the Act and Section 97.5 of the regulations for eligibility requirements.

NOTES: It is the responsibility of the applicant/owner to keep the PVPO informed of any changes of address or change of ownersh or assignment during the life of the application/certificate. There is no charge for filing a change of address. The fee for filing change of ownership or assignment is specified in Section 97.175 of the regulations. (See Section 101 of the Act, and Section 97.130, 97.131, 97.175(h) of Regulations and Rules of Practice.)

To avoid conflict with other variety names in use, the applicant should check the variety names proposed by contacting: See Branch, AMS, USDA, Room 213, Building 306, Beltsville Agricultural Research Center-East, Beltsville, MD 20705. Telephone: (301) 504-8089.

Public reporting burden for this collection of information is estimated to average 30 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Including suggestions for reducing this burden, to Department of Agriculture, Clearant Officer, OIRM, AG Box 7630, Jamie L. Whitten Building, Washington, D.C. 20250. When replying, refer to OMB No 0581-0055 and form number in your letter. Under the PRA of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

The U.S. Department of Agriculture (USDA) prohibits discrimination in its programs on the basis of race, color, national origin, sex, religion, age, dischility, political beliefs, and marital or familie, status. (Not all prohibited bases apply to all programs). Persons with disabilities who require alternative means for communication of program information (braille, large print, audiotape, etc.) should contact the USDA Office of Communications at (202) 720-2791. To file a complaint, write the Secretary of Agriculture, U.S. Department of Agriculture, Washington, D.C. 20250, or call (2027) 720-7327 (voice) or (202) 720-1127 (TDD). USDA is an equal employment opportunity employer.

-52

Attachment for Application Item 20.

	PSU Release ¹	PSU PVP Application ²	PVP Application Date ³	First Commercial Sale
Penn A-1	08-29-95	10-17-96	11-26-96	11-30-95
Penn A-2	08-29-95	10-17-96	12-19-96	02-27-96
Penn A-4	08-29-95	10-17-96	12-19-96	01-24-96
Penn G-1	08-29-95	02-21-97	02-26-97	05-20-96
Penn G-2	08-29-95	10-17-96	12-12-96	01-10-96
Penn G-6	08-29-95	10-17-96	01-02-97	01-24-96
Seaside II	08-29-95	10-17-96	12-12-96	01-26-96

¹PSU Release - Date Pennsylvania Experiment Station Seed Committee approved variety release

²PSU PVP Application - Date on PSU PVP application form

³PVP Application - Application date of variety by PVP office as listed in PVP Official Journal, Vol. 25, December 1997

Origin and Breeding History of Seaside II

The variety Seaside II (experimental designation DF-1) was originally based on a single plant selection from the Desert Forest Country Club in Scottsdale, Arizona in 1984. Selection was made from a segregated patch approximately two feet in diameter on Green No. 8 growing in 15 year old greens originally seeded to Seaside bent. The selection was made based on the segregates density, finer texture, and darker green color compared to the widely segregated greens and determined to be the best performing by then Superintendent Edward Miller and J. M. Duich of Penn State University, the breeder.

Interest in selecting at the site was based on the recognized salt tolerance of Seaside compared to all existing bentgrass cultivars due to the salt content of irrigation water and salt accumulations in the soil under desert climate evaporation conditions. A need exists for salt tolerant bentgrasses with improved turfgrass qualities over Seaside, an ecotype non-bred variety.

The breeding method used was a polycross procedure. The single parent selection of Seaside II was crossed with experimental selections of Penn A and G followed by two generations of crossing selected sib plants of Seaside II.

The original golf course selection was cloned into eight plants, pot planted, and induced to flower in a growth chamber for six weeks. An isolated crossing block was established in the greenhouse in December 1984. Due to known near self sterility of creeping bent, several A and G experimentals were used as pollen parents, the A originated from Penncross greens and G from Penneagle. The A and G plants were fine textured and dense experimentals selected at the August National Golf Club. They were used to introduce a fine leaved texture in crossing. Approximately 250 plants from the crossing were field planted in an isolated planting in the summer of 1985. Some weaker and coarser plants were chemically rogued that fall and reselection continued next year prior to anthesis. Four plants were selected based on vigor, finer density, and uniformity of a darker green color. The four selections followed the procedure used in the breeding program: cloned, growth chamber induced to flower, and crossed with each other to best produce uniformity.

A 300 plant isolated nursery was field planted for reselection in 1987. Fifty plants were selected based on uniformity and similarity to the four selections in the first cycle of reselection and as a convenient number to work with. This nursery appeared more uniform in growth habit and a source of seed for small plot pilot turf testing which ascertained a denser and fine textured turf compared to Seaside. No record now exists of the number of more obvious off-types.

The 50 selections were cloned into three plants and sent to Oregon for evaluation under potential production state conditions. The latter was reduced to 25 clones to serve as potential breeder seed planting stock as the most uniform under Oregon conditions.

Seed from the 25 clones was used to plant a two acre isolated field for evaluation and seed increase. This field was inspected by Oregon Certification and the breeder and considered to be uniform and stable as a variety. Obvious variants were plants identified as more prostrate, more vigorous stoloniferous with lighter green foliage, and estimated to constitute approximately 0.03% of the stand. The uniformity and stability was further ascertained by certification inspection of multiple acreage fields in both the vegetative and reproductive stage.

Commercial production shall be limited to two generations, Breeder and Certified. By agreement with Bent Growers Association with proprietary production rights, all obvious variants are chemically roqued.

Several cycles of five year stands of commercial plantings have shown that Seaside II is a uniform and stable variety to the satisfaction of the breeder and Oregon Certification with no further reselection deemed necessary.

Breeder seed of Seaside II has been maintained and produced under the direction of Pure Seed Testing of Hubbard, Oregon since 1994. Approval of the variety name, Seaside II, has been cleared by the Seed Branch on May 21, 2003.

Variety Distinction of Seaside II

A PVP nursery was established at University Park, Pennsylvania in 1994 consisting of 21 creeping bent varieties with three replications of 25 spaced plants. Included were Seaside II (DF-1), six new Penn State varietal releases, Penn G-1, G-2, G-6, A-1, A-2, A-4, and 13 commercial varieties. Data were collected in 1995 as shown in Table 1. This nursery was discarded due to loss of land.

A second PVP nursery was established at the Pure Seed Testing Research Farm near Hubbard, Oregon in 1995. The purpose was to better evaluate plants in the location of major seed production where growth greatly exceeds the environment in Pennsylvania. It consisted of the above experimentals and 12 commercial varieties with four replications of 25 spaced plants. This test was maintained and data collected and analyzed by Pure Seed personnel after the original PVP application which included only the Pennsylvania location. These data are shown in Tables 2 and 3. Varieties significantly different by years for Seaside II are summarized in Table 4. In this form, the most obvious differences and similarities are easily discerned.

Seaside II may be most easily distinguished from other bentgrass varieties tested by plant height, panicle length, flag leaf length, and base spread. Seaside II differs significantly from Seaside in two or three years for plant height, panicle length, flag leaf width, and number of bottom whorl branches.

In terms of multiple year significance, Seaside II is most similar to Penncross except for plant height.

Table 1. Morphological Character Measurements 1995

	Bottom	Whorl	Branches		5.0	4.7	4.5	4.5	4.5	4.3	4.2	4.2	4.1	4.0	4.0	4.0	4.0	3.9	3.9	3.8	3.8	3.7	3.6	3.6		9.0	
				111111111111111111111111111111111111111	Putter	Seaside	Penncross	Crenshaw	Procup	Penn Â-1	Penn G-2	Penn A-2	Penneagle	Seaside II	Penn G-6	Southshore	Regent	Lopez	Pennlinks	Penn A-4	SR 1020	Cato	Providence	Penn G-1			
	Flag	Leaf	Width	(mm)	2.8	2.7	2.7	2.6	2.6	2.5	2.5	2.5	2.5	2.4	2.4	2.4	2.4	2.3	2.3	2.3	2.2	2.2	2.1	2.1	1.7	0.3	
70000					Seaside	Putter	Regent	Penn A-1	Southshore	Penn A-4	Penneagle	Lopez	Cato	Penn G-1	Penn A-2	SR 1020	Pennlinks	Seaside II	Cobra	Penncross	Penn G-2	Providence	Crenshaw	Penn G-6	Procup		
	Flag	Leaf	Length	(cm)	5.5	5.3	5.2	4.8	4.8	4.7	4.5	4.5	4.4	4.4	4.4	4.3	4.3	4.3	4.2	4.1	4.1	4.1	4.0	3.6		0.5	
77 (247) 11.					Seaside	Penncross	Seaside II	Penn A-1	Penn A-2	Procup	SR 1020	Penn G-2	Penneagle	Pennlinks	Southshore	Penn A-4	Regent	Providence	Penn G-1	Putter	Crenshaw	Lopez	Cato	Penn G-6			
		Panicle	Length	(cm)	8.5	8.3	7.8	7.8	7.8	9.7	7.2	7.1	7.0	6.9	6.9	9.9	9.9	9.9	9.9	6.5	6.4	6.2	0.9	5.7	5.5	9.0	
					Seaside	Cobra	Seaside II	Penncross	Penneagle	Southshore	Regent	Pennlinks	Procup	Putter	SR 1020	Penn A-4	Lopez	Providence	Penn A-1	Cato	Crenshaw	Penn G-2	Penn G-1	Penn G-6	Penn A-2		
-		Base	Spread	(cm)	32.0	33.0	34.0	35.0	36.0	36.0	41.0	44.0	45.0	47.0	47.0	48.0	51.0	55.0	62.0	64.0	65.0	0.69	72.0	80.0	82.0	9.1	
74.65			7,7,7,11.1.1.1	,	Penn G-1	Penn A-1	Penn A-4	Penn A-2	Penn G-2	Penn G-6	Pennlinks	Cato	Crenshaw	Providence	Lopez	Penneagle	Procup	Putter	Regent	Southshore	Cobra	SR 1020	Seaside II	Penncross	Seaside		
	-	Plant	Height	(cm)	40.2	37.8	36.0	34.3	34.3	34.1	34.0	33.7	33.2	33.1	32.8	32.8	31.5	31.1	30.7	30.3	29.8	29.7	29.5	28.4		2.1	
					Seaside	Seaside II	Penn G-1	Penncross	Penn G-2	Penneagle	Pennlinks	Penn A-4	Putter	Southshore	Regent	SR 1020	Crenshaw	Procup	Lopez	Cato	Providence	Penn A-1	Penn A-2	Penn G-6		LSD(0.05)	

¹Penn State University Breeding Nursery, University Park, PA. Three replications of 25 space plants each.

Table 2. Morphological Character Measurements¹ 1996

7.11		•		1111		7447			#
	,				Flag		Flag		Bottom
	Plant		Panicle		Leaf		Leaf		Whorl
	Height	1100000	Length	, ,	Length		Width		Branches
	(cm)		(cm)		(cm)		(mm)		
Seaside II	63.8	Penn A-1	13.1	Penn G-1	9.25	Providence	4.80	Penn A-1	7.3
Seaside	63.1	Seaside	13.0	Seaside II	9.03	Seaside II	4.57	Penn A-2	7.0
Penn A-1	62.5	Southshore	12.8	Seaside	8.90	Lopez	4.53	- Procup	7.0
Southshore	60.2	Crenshaw	12.3	Regent	8.47	Crenshaw	4.52	Crensĥaw	6.7
Penn G-2	57.1	Cato	12.2	Procup	8.26	Penn A-1	4.45	Lopez	9.9
Crenshaw	54.1	Penn G-1	11.7	Penn A-1	7.98	Cato	4.40	Putter	6.4
Providence	53.1	Penn G-6	11.7	Putter	7.91	Procup	4.37	Penneagle	6.4
Penn G-1	53.0	Seaside II	11.6	Penn G-6	7.82	Penncross	4.32	Pennlinks	6.4
Penn A-2	52.2	Penn A-4	11.6	Cato	7.78	Penn G-6	4.17	Penn G-6	6.3
Regent	51.4	Penn G-2	11.4	Penn A-2	7.73	Southshore	4.17	Penncross	6.3
Putter	49.9	Putter	11.2	Crenshaw	7.48	Penn G-2	3.95	Providence	6.3
Lopez	47.5	Pennlinks	11.0	Penn G-2	7.43	Pennlinks	3.83	SR 1020	0.9
Penn A-4	46.7	Regent	11.0	Lopez	7.13	SR 1020	3.82	Cato	5.8
Penneagle	46.0	Penneagle	11.0	Penncross	7.06	Penn A-4	3.77	Penn G-2	5.6
SR 1020	44.3	Procup	10.8	Southshore	7.06	Putter	3.75	Regent	5.6
Procup	44.3	Lopez	10.5	Penneagle	6.87	Penneagle	3.62	Penn G-1	5.6
Penn G-6	43.2	Providence	10.2	SR 1020	98.9	Seaside	3.43	Southshore	5.5
Penncross	43.2	SR 1020	10.0	Providence	89.9	Penn G-1	2.83	Seaside II	5.4
Cato	42.3	Penncross	10.0	Pennlinks	6.50	Penn A-2	2.58	Seaside	4.8
Pennlinks	37.8	Penn A-2	6.6	Penn A-4	5.90	Regent	2.38	Penn A-4	4.6
LSD(0.05)	5.2		0.67		1.56		96.0		0.93

¹Pure Seed Testing Research Farm, Hubbard, Oregon. Four replications of 25 space plants each.

Table 3. Morphological Character Measurements¹ 1997

									#
	ב ק				Flag		Flag		Bottom
	Flami		Panicle T		Leaf		Leaf		Whorl
	Heignt		Length		Length	1,1	Width		Branches
Seaside	(EE)	Seaside	(cm)	Sec.	(cm)	0,000	(mm)	f	1
Penneagle	50.5 40.4	Dennlinke	17.7	December	7.05	Seaside	4.50	Penncross	5.18
Dental Control	7.7.7	reminiks	 	Frovidence	.05	Penneagle	4.27	Penneagle	4.95
renn G-1	6.00	Penneagle	II.	Southshore	7.00	Procup	3.35	Putter	4.95
Seaside III	54.6	Crenshaw	10.7	Penn G-2	6.98	Penncross	3.30	Penn G-1	4.63
Pennlinks	53.9	Providence	10.4	Seaside II	6.97	Lopez	3.23	Seaside II	4.62
Lopez	53.6	Seaside II	10.3	Pennlinks	6.72	Penn G-1	3.12	Lopez	4.53
Penn A-4	53.5	Southshore	10.1	Crenshaw	89.9	Penn A-1	3.00	Penn A-2	4.50
Penn A-2	53.4	SR 1020	10.0	Cato	6.55	Putter	2.95	Providence	4.67
Penn A-1	53.0	Regent	10.0	SR 1020	6.40	Penn A-4	2.93	Southshore	4.42
Penn G-2	52.7	Cato	10.0	Penn G-1	6.04	Pennlinks	2.82	Crenshaw	4.37
Southshore	52.2	Penncross	6.6	Penneagle	5.77	Penn A-2	2.82	Pennlinks	4.35
Regent	51.6	Penn G-1	8.6	Penncross	5.71	Penn G-2	2.73	SR 1020	4.28
Providence	50.1	Penn G-2	9.6	Penn A-1	5.26	Penn G-6	2.70	Regent	4.17
SR 1020	49.8	Penn A-2	9.4	Procup	4.87	Cato	2.70	Seaside	4.17
Cato	49.4	Penn A-1	9.3	Penn A-4	4.82	Providence	2.67	Penn G-2	4.17
Putter	49.1	Putter	9.1	Penn A-2	4.55	Regent	2.58	Cato	4.00
Penn G-6	47.6	Lopez	9.1	Lopez	4.35	Crenshaw	2.58	Penn A-4	3.82
Penncross	47.4	Penn A-4	8.9	Penn G-6	4.29	Seaside II	2.45	Penn A-1	3.60
Procup	46.8	Procup	8.1	Regent	3.97	SR 1020	2.40	Penn G-6	3.45
Crenshaw	46.3	Penn G-6	7.9	Putter	3.83	Southshore	2.37	Procup	3.43
LSD(0.05)	2.9		0.75		0.54		0:30		0.59

¹Pure Seed Testing Research Farm, Hubbard, Oregon. Second Year Test.

Table 4. Varieties Significantly different from Seaside II for Listed Years 1995, 1996, 1997.

read Francis Length Length Width Branches Years 95 96 97 97 97 96 97 90 97 90	110,50
96 97 96 97 96 97 97 96 97 96 97 97 96 97 96 97 96 97 97 97 96 97 96 97 97 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 <t< td=""><td>Flant Height Base Spread</td></t<>	Flant Height Base Spread
96 97 97 97 90 91 96 95 97 96 97 97 97 96 95 96 97 96 97 97 96 97 96 97 96 97 96 96 97 96 97 96 97 96 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 96 97 96 97 96	96
96 95 97 96 97 97 96 97 97 97 96 97 96 97 97 96 97 96 97 96 90 96 97 96 97 97 95 96 97 96 96 97 95 96 97 96 97 96 97 96 97 96 97 95 96 96 96 96 96 96 97 95 96 97 96 97 96 97 96 96 97 96 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96	96
96 96 97 97 97 96 97 96 97 95 97 96 97 96 97 96 97 96 97 95 96 97 96 97 96 97 90 95 96 97 96 97 96 97 96 97 96 95 96 97 96 96 96 96 96 96 96 96 96 96 96 96 96 96 96 96 97 97 97	96 96
96 97 96 97 95 96 97 95 97 96 97 96 97 96 97 96 97 97 95 96 97 96 97 96 97 96 97 95 96 97 96 97 96 96 97 96 97 96 96 97	96 96
96 97 96 97 95 97 97 96 96 97 96 97 96 97 97 95 96 97 96 96 96 95 97 96 97 96 95 96 97 96 97 96 95 96 96 96 96 96 96 97 95 96 96 97 96 96 97 96 97 95 96 97 96	95 96 97 95
96 96 97 96 <td< td=""><td>95 97 95</td></td<>	95 97 95
97 95 96 97 96 97 96 97 96 97 96 96 97 96 96 96 97 96 96 97 96 97 97 96 97 96 96 96 96 96 96 96 96 96 96 96 96 97<	76 96 56
97 96 97 97 96 97 96 97 97 96 97 97 97 97 97 97 97 96 97 96 96 96 96 96 96 96 96 96 96 96 97 96 <td< td=""><td>95 96 97 95</td></td<>	95 96 97 95
96 95 97 97 96 95 96 95 96 97 95 96 96 96 97 97 96 96 96 96 97 97 97 97 97 96 97 97 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 97 96 97 97 97 97 97 97 97 97 97 97	95 96 95
96 95 96 95 96 96 97 95 96 96 96 96 96 96 96 96 96 96 96 96 96 96 96 96 97 96 <td< td=""><td>95 96 97 95</td></td<>	95 96 97 95
96 95 96 96 96 96 95 96 96 96 96 97 85 97 97 97 96 97 96 97 96 97 96 97 96 97 96 97 96 95 96 97 96 97 96 95 96 97 96 97 39 35 22 14 98	95 97
96 95 96 96 96 96 96 96 96 96 96 97 60 97 97 96 <td< td=""><td>95 96 97 95</td></td<>	95 96 97 95
96 95 97 97 95 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 96 97 <td< td=""><td>76 96 56</td></td<>	76 96 56
96 97 97 95 97 96 97 96 97 96 97 96 97 96 96 97 96 96 97 96 96 97 96 96 97 96 96 97 96 96 97 96 96 97 96 97 96 97 96 97 96 97 96 97 <td< td=""><td>95 96 97 95</td></td<>	95 96 97 95
96 97 96 97 96 1 96 95 96 96 1 95 95 95 14 14	95 96 97 95
96 95 96 95 95 39 35 22 14	95 96 95
39 35 22 14	96 97 95
35 22	95 96 97 95
-	46 8 16

U.S. Department of Agriculture Agricultural Marketing Service Science Division Beltsville, Maryland 20705

W-9700058

OBJECTIVE DESCRIPTION OF VARIETY BENTGRASS (Agrostis spp.)

Name of Applicant(s) Pennsylvania Agricultural Experiment Station	Variety Name or Temporary
	Designation
Address (Street and No. or R.F.D. No, City, & ZIP Code)	Seaside II (DF-1)
0217 Agricultural Administration Bldg.	FOR OFFICIAL USE ONLY
The Pennsylvania State University	PVPO Number
University Park, PA 16802	<u> </u>
Place numbers in the boxes (e.g. 089) for the typical plants of this variety. The sym	e characters that best describe bol & indicates decimal.
COMPARISON VARIETIES FOR USE BELOW	
l= Astoria 2= Exeter 3= Highland 4= Seaside 7= Astra 6= Other Penneagle	5= Penncross 6= Kingstown
	eping A. stolonifers (A.palustr wn bent A. canina ssp. montana
2. ADAPTATION: (0= Not Tested, 1= Not Adapted,	·
Northeast Southeast North Central Other (Specify) South West	_
Northeast Southeast North Central Other (Specify) South West 3. MATURITY (At first anthesis): Use comparison	-
2 Other (Specify) South West	varieties
2 Other (Specify) South West 3. MATURITY (At first enthesis): Use comparison 0 4 Days earlier than 5 , Maturity same as 4	varieties ,0 8 Days later than 8
2 Other (Specify) South West 3. MATURITY (At first anthesis): Use comparison 0 4 Days earlier than 5 , Maturity same as 4 4. HEIGHT (Average of longest 10 shoots from soil	varieties , 0 8 Days later than 8 later than 1 surface to top of head):
2 Other (Specify) South West 3. MATURITY (At first anthesis): Use comparison 0 4 Days earlier than 5 , Maturity same as 4 4. HEIGHT (Average of longest 10 shoots from soil	varieties ,0 8 Days later than 8 later than 4 later than 8 later than
2 Other (Specify) South West 3. MATURITY (At first anthesis): Use comparison 0 4 Days earlier than 5 , Maturity same as 4 4. HEIGHT (Average of longest 10 shoots from soil 5 2 Cm Height (at maturity) Ave 0 3 Cm Short	varieties ,0 8Days later than 8 l surface to top of head): rter than 4 same as Comparison
2 Other (Specify) South West 3. MATURITY (At first anthesis): Use comparison 0 4 Days earlier than 5 , Maturity same as 4 4. HEIGHT (Average of longest 10 shoots from soil 5 2 Cm Height (at maturity) Ave 0 3 Cm Shoots Height	varieties ,0 8Days later than 8; l surface to top of head): rter than 4 Comparison
2 Other (Specify) South West 3. MATURITY (At first anthesis): Use comparison 0 4 Days earlier than 5 , Maturity same as 4 4. HEIGHT (Average of longest 10 shoots from soil 5 2 Cm Height (at maturity) Ave 0 3 Cm Shoots Height 10 Cm Tal	varieties , 0 8 Days later than 8 lead lead lead lead lead lead lead lead
2 Other (Specify) South West 3. MATURITY (At first anthesis): Use comparison 0 4 Days earlier than 5 , Maturity same as 4 4. HEIGHT (Average of longest 10 shoots from soil 5 2 Cm Height (at maturity) Ave 0 3 Cm Shoots Height	varieties , 0 8 Days later than 8 lead lead lead lead lead lead lead lead

Bentgrass - 2 -

6. VEGETATIVE REPRODUCTION:	· · · · · · · · · · · · · · · · · · ·
Rhizomes l= Absent 2= Present	2 Stolons 1= Absent 2= Present
Rhizcmes 1100 & Stolons	PAD
7. LEAF BLADE:	8/9/0
3 Color: 1- Yellowish Green (Cohansey) 3- Green (Exeter)	2= Light Green (Washington)
5= Bluish Green (Highland)	h= Dark Green (Kingstown, Tracenta 6= Other (Specify)
2 Texture: (fineness) 1- Very fine (Kingstown) 3- Medium fine (Astoria) 5- Medium coarse (Virginia)	2= Fine (Exeter) 4= Medium (Seaside) 6= Coarse (Vermont)
Stomatal density upper leaf surface (Num	mber/mm ²)
Lower Surface:]≴ Rough
Upper Surface: Smooth	% Rough
Margins: \$ Smooth []	% Rough
Mm Width (Amount	arrower than []
	comparison
-	der than
3.11 Mr. 175-343 (77)	ngth (Flag leaves)
8. LEAF SHEATH:	
Anthocyanin: 1= Absent 2= Present	% Red sheaths
9. LIGUIE (Lower and middle leaves):	
Shape at Apex: 01610 % Acute 7 Ro	ounded 02 400 % Trupcate
5 Other (Specify)	- I wicave
Pubescence: 11010 Glabrous	% Pube scent
Margins: 0 30 % Entire 070 %	Toothed
% Other (Specify)	rootned
Mm Length	

10. LEMMA:
Shape: Sh
3 9 Mm Width 115 Mm Length (exclusive of awn)
Color: Buff 100 % Silvery Cher (Specify)
Surface 100 % Glossy Dull
Texture: 100% Smooth % Punctate
Pubescence:
Basal Hairs: 100 % Absent Few Amy
Short I S Long
Appressed
Awns: 100 % Absent Few Many
% Awn-pointed % Short M % Long
Straight Geniculate
Awn Insertion on Lemma:
% Basal
11. PANICLE:
Type (in anthesis): 0 4 0 % Open 0 6 0 % Compact
Anthocyanin: 0 70 % Absent 0 3 0 % Present
Branches in Anthesis: 0 70 % Appressed 0 3 0 % Ascending 7 % Spreadi
Branches in Fruit: 0.50 % Appressed 050 % Ascending 7 Spreadi
Branch Surface: 070 & Smooth 030 & Scabrous
12. SEED:
8 8 Frams per 1000 seed
13. SPRING GREEN UP:
2 l= Early (Exeter) 2= Medium (Astoria) 3= Late (Kingstown)

Bentgrass - 4 -

14. ENVIRONMENT	LL RESISTANCE: (0=)	Not te	sted, l= Suscepti	ble 2= Resistant	:1
	Heat Drough	_		er (Specify)	·)
15. DISEASE RESI	STANCE (O= Not te:	sted	l= Susceptible 2=	Resistant):	
2 Red Leaf Sp	ot - Drechslera erg - Drechslera poae (Helminthosporiu	thros	pila Helmintho (Bipo)		~ \
Pythium Bli	ght - (P. aphanider	matum)	Pythium B	light (P. ultimu	n)
Fusarium Bl:	ight (F. roseum)			Blight (E tricing	
(F_nival	ch (Pink Snow Mold			ldew (Erysiphe	
1 Ophiobolus F	etch (O. graminis)		Stripe Smu	t (<u>Ustilago str</u>	
_	(Gloeocercospora so		(T. in	ight (Snow Scald	
	Corticium fuciformuccinia graminis)	<u>ne</u>)	Brown Patc	h (Rhizoctonia	solani)
	• pose-nemoralis)		Crown Rust	(P. coronata)	
					
	NCE (> Not tested	, 1= 5	Susceptible, 2= Re	esistant):	
European Chai	Cer Control		Garden Chaf	·	
(Amphimallo	n solstitialis)				
Chinch Bug (E	lissus insularis)		Webworm (Cr	ertha horticola)
Armyworm (Cut	warm)				
(Pseudalet	ia unipuncta)		1 Other Blac	k Cutworm	·
<u> </u>		· •· •	·		
following number) THAT MOST CLOSELY acteristics indicaters; 1= Submitted y, 2= Same as, 3=	mond of	ree of resemblance	e (D.R.) with on	the e of the rior to
haracter	.,	*******	chan, darker or	superior, etc.	
rowth Habit	Similar Variety Penncross	D.R.	Character	Similar Variety	D.R.
wn Length	. remicross	2	Leaf Color	Penncross	2
eed Weight	<u> </u>		Panicle Type	Penncross	2
old Resistance	Penncross	-	Turf Fineness	Penncross	3
rought Resistance	, , remicross	2	Heat Resistance		
rown Patch	Penncross	3 -	Shade Resistance		
ensity	Penncross		Moss Resistance		
3. COMMENTS:			Salt Tolerance	Penncross	3

Additional Description of Seaside II

The primary objective of the breeding program was to improve the turfgrass quality of Seaside II over Seaside but to retain salt tolerance. A salt tolerance test in greenhouse sand culture was conducted at Pure Seed Testing in Hubbard, Oregon in a salt concentration of 3000 ppm of NaCl and 5% CaCl. In this heavy salt regime 55% of Seaside II plants survived vs 64% of Seaside. All other bentgrasses in the 1996 PVP test showed 0 to 2% survival, the latter with poor growth. Significant improvement in turfgrass differences versus Seaside with the breeding and selection porcedures are as follows.

Table 5	Leaf texture: 0.74 vs1.01, 0.76 vs 0.99, 0.99 vs 0.90
Table 6	Leaf texture: 6.8 vs 4.6 rating, 1995 NTEP
Table 7	Genetic color: 6.0 vs 4.9 rating, 1994 NTEP
Table 8	Shoot density: 2058 vs 765
Table 9	Summer density: 7.3 vs 4.4 rating, 1995 NTEP
Table 10	Turf quality: 6.0 vs 4.5, 1995 NTEP
Table 11	Brownpatch rating: 7.6 vs 6.1, 1994 NTEP
Table 12	Moss percent: 3.8 vs 31.7% in 1993; 7.4 vs 26.8% in 1994

Table 5. Leaf texture of putting green bent maintained at 4.0 mm as putting green turf in three locations.

University	ity Park, PA		Augusta, GA	a, GA	Turin, Italy	[taly
	1993	1999		1993		
Penn G-2	0.61	0.73	Penn A-2	0.63	Penn G-2	0.63
Penn G-6	0.63	0.73	Penn A-1	0.65	Penn G-6	0.70
Penn A-1	0.63	0.75	Penn G-2	0.65	Penn A-1	0.70
FHG-1	0.63		Penn G-1	0.68	Penn G-1	0.72
Penn G-1	0.65	69.0	Penn A-4	69.0	Seaside II	0.79
Penn A-2	29.0	0.71	Penn G-6	0.71	Pennlinks	0.80
Penn A-4	0.69	0.76	Crenshaw	0.79	SR-1020	0.84
Seaside II	0.74	0.76	Cato	0.80	Southshore	0.85
Pennlinks	0.77	0.93	Seaside II	0.80	Penncross	0,85
Cato	0.80	0.84	Penncross	0.99	Providence	0.86
SR-1020	0.80	0.84			Putter	0.88
Providence	0.81	96.0			Cobra	0.90
Penneagle	0.85	0.95			National	06.0
Putter	0.89				Seaside	0.90
Carmen	0.93				Penneagle	0.95
Cobra	0.95				Emerald	96.0
Penncross	0.99	0.99				
Seaside	1.01	0.99				
Emerald	1.12					
LSD (0.05)	0.04	0.09		0.06		0.10

¹Leaf width of second sub-tended leaf (mm).

					LEAF	TEXTURE	LEAF TEXTURE RATINGS 1-9; 9=VERY	3S: 1-9;	9=VERY	FINE							
NAME	141	1.1	LNI LNI	KS2	KX	MB.1	5	MI	M02	2	H	7.	ua1	VAB	WA1	WA3	MEAN
PENN G-2 (G-2)	7.0	8.7	7.0	8.3	0.6	6.3	7.7	7	K	c	,	c	6	•	•	1	. ,
PENN G-6 (G-6)	7.0	0.6	7.3	8.7	8.0	7	~	1	, ,		- 6	2 (0 0		⊃ : o :	5.	7.6
BAR WS 42102	7.7	8	8	0			, ,		- 1) c	0.0) ·	ۍ د د	• •	7.3	7.3	٠.
CATO	~	7	2 2) ! 	3;	ə !	ا c	6.3	œ. 0.	9.0	ω Υ.	7.7	7.2
STAITHGHODE			- 1	0 6	- (6.5	2.7	7.7	6. 7	۷.	6.7	8.0	5.0	8.0	7.0	2.0
	- 1	> 0		, r	ع د د د	0.4	0. 0	0.9	7.7	6.3	8	6.0	7.7	5.0	7.7	7.3	7.0
ACTION TO THE PERSON AND THE PERSON	- !) (3,	 	(·	0.4	7.3	۲.	7.3	5.7	8.0	2.7	8.0	4.7	8.0	6.3	6.9
SERSINE LI (UF-1)	1	0.4	0.7	8.3	8.3	3.7	2.0	9.0	7.0	4.3	8.0	7.0	0.8	5,0	7.0	2.0	
PROVIDENCE	7.3	œ .	7.3	8.0	7.0	3.7	6.3	5	akai kai	9.0	7.7	5.3	6	4.7			2
PENNEAGLE	7.3	œ.	7.3	8.7	7.0	3.7	5.7	0.9	7.0	5.0	0.0	9.0	8	0.4	7.7	2	4
SK 7100	6.3	0.0	7.0	2.7	7.0	2.3	6,3	ν. Μ	8.0	6.0	8.0	6.0	7.7	N.		~	· ·
1S1-AT-90162	6.3	ب س ا	6.7	5.3	6.7	3.7	6.0	5.0	8.7	5.0	7.0	6.3	0.8	0	7.0		. 4
18TH GREEN	0 ·	7 7	7.0	7.3	8.0	0.4	5.7	4.7	7.7	5.0	8.0	5.0	7.0		7.7		7
OM-AT-90163	2 /	۳ ا	7.0	7.7	2.7	4.0	5.7	5.0	8.3	5.0	7.3	7.0	8.0	5.0	0.7	9	· ·
LOPEZ	2 1	7.7	7.0	7.3	7.7	M M	6. 0	5.7	7.3	4.3	0.8	6.0	7.0	P.	2.0	ır.	, ,
TRUELINE	7.3	7.7	7.0	7.0	6.3	0.4	6.0	5.0	7.0	3.3	7.7	ν,	7.0	7	2	. 4	
PRO/CUP	9	7.7	6.7	7.0	7.0	3.3	5,3	5.3	7.0	3.7	0.8	5.7	7.0	7	2.7	, r.	
BAR AS 492	5.3	8.3	7.0	5.7	2.0	7.2	6.0	4.3	8.0	5.0	7.3	6.3	7.7	7.7	4		
PENNCROSS	8	7.3	7.3	6.3	2.9	4.3	6.0	5. 0.	7.3	2.3	7.3	6.0	7.0	7	W V		, c
TENDENZ	S.	8.7	6.7	4.3	7.0	3.0	5.7	5.0	7.7	4.3	7.7	7.7	7.7		1 4	. 7	
EXETER	7.7	7.7	6.7	7.7	6.0	2.0	5.7	5.0	8.3	3.0	7.7	6.3	. M	, M	4	۳ س	, n
SEASIDE	6.3	2.0	5.3	3.7	5.0	2.3	5.0	4.0	7.0	1.0	7.7	4.7	6.0	2.7	4.3	2.3	4.6
LSD VALUE	1.9	7.0	1.0	0.9	0.9	1.3	6.0	0.6	0.8	1.2	0.7	1.5	0.5	-:	0.9	1.2	0.3

TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05). **-**

GENETIC COLOR RATINGS OF BENTGRASS CULTIVARS
GROWN ON A FAIRWAY OR TEE
1994 DATA

7
GREEN
9=DARK
4-0
RATINGS
COLOR
GENET 1C

										:					
NAME	1-1	KS2	KX 1	MA.	MB 1	E E	OH1	PA1	R11	UB1	VA8	WA1	HA3	NI 1	MEAN
18TH GREEN	7.3	8.0	8.3	8.3	9.0	7.0	7.0	8.0	7.0	0,8	0.5	۲,	7.0	۲ ۲	7 5
CATO	7.7	8.3	7.7	8.0	7.7	7.0	7.3	5.	6.7	7.3	. C		W .	,	
TENDENZ	ν. 	8.0	8.3	7.3	7.3	7.0	8.0	8.7	7.0	6.0	6.7	7		C	2
PROVIDENCE.	7.7	7.3	7.7	8.0	7.0	8.0	7.0	6.7	5.7	7.0	0.0	0.0	5.7	9	9
CRENSHAW	7.3	7.0	8.0	8.3	6.7	6.7	6.7	7.7	6.0	7.0	5.0	9.0	5.0	6.7	6.9
1S1-AT-90162	5.3	7.7	8.0	6.3	6.3	7.0	8.0	7.0	4.7	0.9	6.3	6.0	7.0	0.8	6.7
9-9	7.3	8.0	7.7	8.0	6.7	7.0	7.3	4.0	5,3	8.0	5.0	8.0	4.3	6.7	6.7
OM-AT-90163	6.0	7.7	7.7	6.7	7.3	7.0	7.7	7.0	4.0	6.0	2.9	6.0	6.0	7.7	6.7
G-2	7.0	8.0	7.3	8.0	6.7	6.0	7.0	6.0	6.0	7.7	5.7	8.0	.3	6.9	9.9
LOPEZ	6.7	7.3	6.7	6.7	7.3	7.7	7.3	5.7	6.3	6.3	5.0	7.7	6.0	6.9	9.9
SOUTHSHORE	7.3	7.0	8 .0	5.7	6.7	6.0	7.0	7.0	5.0	7.0	5.0	8.0	5.0	6.0	6.5
BAR WS 42102	7.3	8.0	8. 0.	6.7	6.7	7.0	7.0	4.0	5.3	7.0	5.0	8.0	4.7	5.4	6.5
TRUEL INE	6.7	7.3	6.3	7.0	7.0	6.3	2.0	5.0	6.0	7.0	5.0	7.7	6.0	6.0	6.5
PRO/CUP	2.9	7.0	8.0	6.7	6.7	7.0	7.0	5.7	5.0	6.3	5.0	7.7	5.0	6.0	7.9
PENNEAGLE	7.3	7.0	7.3	6.7	6.7	7.0	7.0	5.3	5.0	7.0	5.0	7.3	4.0	6.0	6.3
PENNCROSS	7.0	7.0	7.3	7.0	9.0	7.0	7.3	5.7	5.0	6.3	4.0	7.3	5.3	6.3	6.3
SR 7100	5.7	7.0	7.7	6.3	6.7	7.0	7.3	6.0	3.7	6.0	6.3	5,3	5.0	8.0	6.3
BAR AS 492	6.3	6.3	6.3	6.0	7.0	7.0	7.3	5.0	4.0	6.0	6.3	6.3	4.3	6.7	6.1
DF-1	7.0	6.7	7.0	7.0	6.0	6.0	7.0	4.0	5.0	6.7	5.0	7.3	2.7	6.0	6,0
EXETER	7.0	2.7	6.0	4.0	6.7	7.0	6.0	4.3	4.7	5.0	5.3	9.9	5.3	6.3	5.7
SEASIDE	6.0	5.0	6.3	4.0	6.0	5.3	7.0	2.0	3.0	5.0	5.7	5.7	1.7	6.0	4.9
LSD VALUE	1.0	6.0	0.9	1.3	6.0	9.0	7.0	1.2	6.0	0.5	9.0	6.0	1.2	0.7	0.2

TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05). 2

Table 8 Bentgrass shoot density/dm², Augusta, GA, and Turin, Italy.

	Augus	sta, GA	_	Turi	n, Italy
	1992	1993	_	<u>1992</u>	1993
PENN A-2 PENN A-1 PENN G-1 PENN G-2 PENN A-4 PENN G-6 Crenshaw Cato Penncross Seaside II	2376 1815 1881 2079 1617 1683 1617 1254 1122 1419	2392 2145 1996 1963 1917 1838 1419 1287 1270 1056	PENN G-1 PENN G-2 PENN G-6 PENN A-1 Seaside II Southshore Pennlinks Providence SR 1020 Putter Penneagle Cobra National Emerald Seaside	1574 1080 1065 1075 1043 1000 914 1017 1091 980 1170 908 915 591	2612 2546 2378 2240 2058 1509 1504 1425 1419 1272 1241 1196 1013 1010 765
LSD (.05)	180	214	Comitac	258	178

SUMMER DENSITY RATINGS OF BENTGRASS CULTIVARS GROWN ON A FAIRWAY OR TEE 1995 DATA

1
DENSITY
9=MAXIMUM DE
1-9;
RATINGS
DENSITY

NАМЕ	38	PA1	HA3	MEAN
PROVIDENCE	8.0	8.0	7.7	7 0
PENN G-2 (G-2)	8.7	8.0	7.0	0.2
18TH GREEN	9.0	6.7	7.0	2.6
BAR WS 42102	8.0	7.3	7.3	7.6
CRENSHAW	8.7	7.0	6.7	7.4
PENN G-6 (G-6)	8.0	7.7	6.3	7.3
SEASIDE II (DF-1)	8.3	8,3	73	7.3
SOUTHSHORE	1.7	-	ر و	<u> </u>
TRUELINE	8.3	6.7	6.7	7.2
CATO	7.7	7.0	6.3	7.0
PENNEAGLE	8,3	7.0	5.7	2.0
PRO/CUP	8.3	5.7	6.7	6.9
ISI-AT-90162	7.3	6.9	7.0	6.8
LOPEZ	7.7	6.3	6.3	
BAR AS 492	7.3	6.7	2 2	
PENNCROSS	8.3		6.0	
SR 7100	7.0	5.7	5.0	
TENDENZ	7.7	5,3	7	
OM-AT-90163	7.3	5.3	3.7	5.4
EXETER	6.0	4.7	7	
SEASIDE	6.7	3.3	3.3	4.4
LSD VALUE	1.1	-	7	
10101	•	:	<u>:</u>	

TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05). **-**

0.7 189 0.7 6.0 PA1 MEAN TURFGRASS QUALITY RATINGS OF BENTGRASS CULTIVARS GROWN ON A FAIRWAY OR TEE AT NINETEEN LOCATIONS IN THE U.S. AND CANADA 8.7.7.8 7.7.7.7.5 7.7.5 7.3 띪 7.0 TURFGRASS QUALITY RATINGS 1-9; 9=1DEAL TURF 9.0 3 7.3 7.2 7.0 7.0 7.0 7.0 7.0 7.0 7.0 7.0 7.0 0.5 吾 1995 DATA 9.0 Ξ € 6.0 9.0 9.0 4.6 5.0 5.0 5.0 5.0 MA1 KY 8.0 7.6 8.0 7.7 9.7 0.7 ₹. Ξ 7.0 4.6 7.3 6.9 6.7 6.4 6.1 6.1 -Σ SEASIDE 11 (DF-1) CRENSHAW PENN G-2 (G-2) PENN G-6 (G-6) ISI-AT-90162 BAR WS 42102 SR 7100 OM-AT-90163 SOUTHSHORE BAR AS 492 18TH GREEN **PROVIDENCE** PEHNEAGLE PENNCROSS TRUEL INE Lopez LSD VALUE PRO/CUP TENDENZ EXETER

လလန္လန္လန္လလ္လ ပ်ပစ္ဝဲဆ်ဆံလ်လ်ပမေ

COMMERCIALLY AVAILABLE IN THE USA IN 1996

TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05). _

BROWN PATCH RATINGS OF BENTGRASS CULTIVARS GROWN ON A FAIRWAY OR TEE 1994 DATA

BROWN PATCH RATINGS 1-9; 9=NO DISEASE 1/

				•			
	NAME	MA1	MO1	M02	1, N	PA1	MEAN
	PROVIDENCE	0.6	7.3	8	8.7		ر در
	05-1	9.0	3.3	0	8.7	0.0	· ~
١.	TRUELINE	8.7	3.3	₽ 8			• 12
	CATO	9.0	3,3	2.7	8	0.0	
	6-2	0.6	2.7	8	8.3	0.0	
	9-9	0.6	2.3	7.7	0.0	0.0	7 /
	18TH GREEN	8.3	2.3	× 8	8.7	0.6	2
	LOPEZ	8.3	4.0	6.7	8.7	0.6	7.3
	SOUTHSHORE	0.6	2.0	8.0	8.3	9.0	7.3
	PRO/CUP	0.6	2.7	7.7	8.0	0.6	7.3
	CRENSHAW	0.6	2.0	6.7	9.0	0.6	7.1
	PENNEAGLE	0.6	2.7	6.3	8.7	0.0	7
	BAR WS 42102	8.3	2.3	7.7	7.3	0.0	9
	SR 7100	5.7	7.3	5.0	6.3	2,3	9
	PENNCROSS	8.3	3.0	5.3	5.7	0	1
	EXETER	8.3	3.7	4.7	4.7	0.0	6.1
	SEASIDE	8.0	2.7	5.7	5.0	0.6	6.1
	ISI-AT-90162	7.3	3.0	5.3	5.0	0.	5.5
	BAR AS 492	7.0	4.7	3.7		6.3	, r.
	TENDENZ	7.3	2.0	3.0	4.0	5.3	6.3
	OH-AT-90163	6.0	2.0	2.7	4.0	6.3	4.2
	1 SP WALLE	9	7	r	,		•
	ראה אשרתר	0	0	۲.5	?	8.0	٠.

TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05). **_**

Table 12 Presence of moss in 1993 and 1994. Turin, Italy.

		··· ·· ·· ·
	Moss-1993 (%)	Moss-1994 (%)
PENN G-1	0.10	1.93
PENN A-1	0.05	1.14
PENN G-6	0.33	3.64
Southshore	0.72	3.67
Penneagle	2.55	5.19
PENN G-2	0.55	2.64
Putter	3.94	6.67
Pennlinks	3.19	7.19
Providence	2.86	3.00
Penncross	3.36	12.29
Seaside II	3.80	7.36
Cobra	3.58	11.19
SR 1020	4.08	8.67
National	11.17	18.81
Emerald	9.33	17.76
Seaside	31.67	26.78
Astoria	24.77	35.57
LSD (.05)	10.20	6.81

REPRODUCE LOCALLY. Include form number and date on all reproductions. U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE	FORM APPROVED - OMB I	
SCIENCE AND TECHNOLOGY DIVISION - PLANT VARIETY PROTECTION OFFICE	1974 (5 U.S.C. 652a) and the Pay	de in eccordance with the Privacy Act perwork Reduction Act (PRA) of 1995.
EXHIBIT E STATEMENT OF THE BASIS OF OWNERSHIP	Application to assure to the con-	o determine if a plant variety protecti
1. NAME OF APPLICANT(S)	2. TEMPORARY DESIGNATION OR EXPERIMENTAL NUMBER	3. VARIETY NAME
Pennsylvania Agricultural Experiment Station	DF-1	Seaside II Creeping Bentgrass
4. ADDRESS (Street and No., or R.F.D. No., City, State, and ZIP Code, and Country)		
Charles R. Krueger Bruce McPheron Associate Dean	5. TELEPHONE (Include area code)	6. FAX (include area code)
0217 Agricultural Admin. Bldg.	814-865-5410 7. PVPO NUMBER	814-863-7905
University Park, PA 16802	9700	1058
8. Does the applicant own all rights to the variety? Mark an "X" in appropriate bi	lock. If no, please explain.	
		X YES NO
		*
	<u> </u>	
 Is the applicant (individual or company) a U.S. national or U.S. based company? If no, give name of country 		X YES NO
O. Is the applicant the original breeder? If no, please answer the following:		
 a. If original rights to variety were owned by individual(s): ls (are) the original breeder(s) a U.S. national(s)? If no, give name of cor 		YES X NO
o and a distribution of the state of the sta	untry	
h If acidian Line and a control of the control of t	Г:	YES NO
 b. If original rights to variety were owned by a company: ls the original breeder(s) U.S. based company? If no, give name of count 	try	AYES NO
11. Additional explantion on ownership (If needed, use reverse for extra space):		
J. M. Duich, Professor Turfgrass Science is breed Employee of applicant with rights assigned to app	er. licant.	•
LEASE NOTE:		
ant variety protection can be afforded only to owners (not licensees) who meet one	of the following criteria:	
If the rights to the variety are owned by the original breeder, that person must be of a country which affords similar protection to nationals of the U.S. for the same		OV member country, or national
If the rights to the variety are owned by the company which employed the original nationals of a UPOV member country, or owned by nationals of a country which affigenus and species.		U.S. based, owned by

3. If the applicant is an owner who is not the original breeder, both the original breeder and the applicant must meet one of the above criteria.

The original breeder may be the individual or company who directed final breeding. See Section 41(a)(2) of the Plant Variety Protection Act for definition.

Public reporting burden for this collection of information is estimated to average 10 minutes per response, including the time for reviewing instructions, searching existing data sources, gethering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including usagestions for reducing this burden, to Department of Agriculture, Clearance Officer, OIRM, AG Box 7630, Jamie L. Whitten Building, Washington, D.C. 20250. When replying, refer to OMB No.

Under the PRA of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

The U.S. Department of Agriculture (USDA) prohibits discrimination in its programs on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, and marital or familial status. (Not all programs). Persons with disabilities who require alternative means for communication of program information (braille, large print, audiotape, etc.) should contact the USDA Office of Communications at (2021) 720-2791.

To file a complaint, write the Socretary of Agriculture, U.S. Department of Agriculture, Washington, D.C. 20250, or call (2021 720-7327 (voice) or (2021 720-1127 (TDD). USDA is an equal employment opportunity employer.